


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




4089
4090
4091
4092
4093
4094
4095
4096
4097
4098
4099
4100
4101
4102
4103
4104
4105
4106

Wilbur H. Henig Campbell. Affinity binding via Zinc(II) for controlled orientation and electrochemistry of Histidine-tagged nitrate reductase in self-assembled monolayers. <i>Bioelectrochemistry</i> . 2013. 93:46-50
L. Albinet Chaal. Wall shear stress mapping in the rotating cage geometry and evaluation of drag reduction efficiency using an electrochemical method. <i>Corrosion Science</i> . 2009. 51:1809-1816
Jorge N. Chedekel Chac. Chemically induced parkinson's disease II: Intermediates in the oxidation and reduction reactions of the 1-methyl-4-phenyl-2,3-dihydropyridinium ion and its deprotonated form. <i>Biochemical and Biophysical Research Communications</i> . 1989. 158:63-71
Surendra Mohanty Chandra Sabat. Characterization of Heat-stress Induced Stimulation of Photosystem I Electron Transport Activity in Amaranthus Chloroplasts: Effect of Cations. <i>Journal of Plant Physiology</i> . 1989. 133:686-691
Wei-Chieh Deka Chang. Synthesis and characterization of large pore cubic mesoporous silicas functionalized with high contents of carboxylic acid groups and their use as adsorbents. <i>Applied Catalysis B: Environmental</i> . 2013. 142:817-827
Neelam P. Fatma Chauhan. Protection of Wheat Chloroplasts from Lipid Peroxidation and Loss of Photosynthetic Pigments by Quercetin under strong Illumination. <i>Journal of Plant Physiology</i> . 1992. 140:409-413
M. E. Lang McManus. Activation of misonidazole by rat liver microsomes and purified NADPH-cytochrome c reductase. <i>Biochem Pharmacol</i> . 1982. 31:547-52
Hao Yang Chen. A fluorescent guest used to determinate the effective content of CB[8] and to further detect methyl viologen. <i>Chinese Chemical Letters</i> . 2013. 24:857-860
Zhen Juneau Chen. Effects of three pesticides on the growth, photosynthesis and photoinhibition of the edible cyanobacterium Ge-Xian-Mi (Nostoc). <i>Aquatic Toxicology</i> . 2007. 81:256-265
Mordechai Chevion. A site-specific mechanism for free radical induced biological damage: The essential role of redox-active transition metals. <i>Free Radical Biology and Medicine</i> . 1988. 5:27-37
Alexander K. Slavnova Chibisov. Electron transfer reactions induced by the triplet state of thiocarbocyanine dimers. <i>Chemical Physics</i> . 2004. 299:10-Jan
Nakul K. Choe Choudhury. Ascorbate induced zeaxanthin formation in wheat leaves and photoprotection of pigment and photochemical activities during aging of chloroplasts in light. <i>Journal of Plant Physiology</i> . 1993. 141:551-556
Nam-Hai Chua. Photooxidation of 3,3'-diaminobenzidine by blue-green algae and Chlamydomonas reinhardtii. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1972. 267:179-189
Lakshaman Kumar Sawhney Chugh. Photosynthetic activities of Pisum sativum seedlings grown in presence of cadmium. <i>Plant Physiology and Biochemistry</i> . 1999. 37:297-303
Prakit Jakmunee Chuntib. Simple flow injection colorimetric system for determination of paraquat in natural water. <i>Talanta</i> . 2015. 144:432-438
Robyn E. Grace Cleland. Voltammetric detection of superoxide production by photosystem II. <i>FEBS Letters</i> . 1999. 457:348-352
Annette Vignais Colbeau. The membrane-bound hydrogenase of Rhodospseudomonas capsulata: Stability and catalytic properties. <i>Biochimica et Biophysica Acta (BBA) - Enzymology</i> . 1981. 662:271-284
Graeme Daniels Cooke. The synthesis of a pyrrole-functionalized cyclobis(paraquat-p-phenylene) derivative and its corresponding [2]rotaxane and [2]catenane and their subsequent deposition onto an electrode surface. <i>Tetrahedron</i> . 2007. 63:11114-11121

Monolayers of CuII-complexes on electrode surfaces are frequently applied for the immobilization and controlled orientation of biomolecules.
The local wall shear stress (WSS) mapping in the rotating cage (RC) has been obtained from measuring the diffusion current of a redox probe.
The one-electron reduction product of 1-methyl-4-phenyl-2,3-dihydropyridinium ion has been generated by pulse radiolysis and characterized by laser Raman spectroscopy.
Summary Exposure of isolated Amaranthus chloroplasts to elevated temperatures (>25 °C) results in a rapid loss of photosynthetic activity.
Well-ordered cubic FDU-12 type mesoporous silicas functionalized with various contents of carboxylic acid group (COOH) have been synthesized by a sol-gel process.
Summary Exposure of chloroplasts to strong visible light in the presence of DCMU and paraquat resulted in lipid peroxidation and release of superoxide anion.
Rat liver microsomes and purified NADPH-cytochrome c reductase metabolized [14C]misonidazole anaerobically to a range of products.
A fluorescent pyrene derivate, N-allyl-1-pyrenemethylammonium hydrochloride (APA+), was reported to form a stable helical dimer in aqueous solution.
Effects of butachlor, bensulfuron-methyl, and dimethoate on the growth, photosynthesis, and photoinhibition of the edible bean (Phaseolus vulgaris L.) were studied.
The metal-mediated site-specific mechanism for free radical-induced biological damage is reviewed. According to this mechanism, the metal ion acts as a catalyst for the generation of free radicals.
The photoinduced electron transfer between either cationic 5,5'-dichloro-3,3',9-triethylthiacarbocyanine (1) or a structurally related dye and a series of electron acceptors was studied.
Summary Mechanisms by which higher levels of Zeaxanthin (Zx) in detached wheat leaves, induced by ascorbate in vivo, are related to the activation of the xanthophyll cycle.
1. 1. The photooxidation of 3,3'-diaminobenzidine was investigated in whole cells of the wild-type and two mutant strains of Escherichia coli.
One-month-old pea seedlings (Pisum sativum L. cv. Bonneville) raised in sand culture, were provided with a nutrient solution containing different concentrations of paraquat.
A simple and low cost flow injection colorimetric system has been developed for determination of paraquat in natural water samples.
Oxygen radicals play both pathological and physiological roles in biological systems. The detection of such radicals is difficult and the development of sensitive and specific probes is a major challenge.
The hydrogenase of Rhodospseudomonas capsulata is an intrinsic membrane protein extractable from the membrane by treatment with detergents.
We report the convenient synthesis of a pyrrole-functionalized tetracationic cyclophane, [2]rotaxane, and [2]catenane. X-ray crystallographic analysis of the [2]rotaxane shows that the pyrrole ring is located inside the cyclophane cavity.

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



4107
4108
4109
4110
4111
4112
4113
4114
4115
4116
4117
4118
4119
4120
4121
4122
4123
4124
4125

Sylvie Cournac Corneille. Reduction of the plastoquinone pool by exogenous NADH and NADPH in higher plant chloroplasts: Characterization of a NAD(P)H. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1998. 1363:59-69
S. Da Silva Cosnier. Electrochemical nitrate biosensor based on poly(pyrrole). <i>Bioelectrochemistry</i> . 2008. 74:47-51
Sean J. Coughlan. Chloroplast thylakoid protein phosphorylation is influenced by mutations in the cytochrome bf complex. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1988. 933:413-422
Paulo J. G. Costa Coutinho. Kinetics of the electron transfer reaction between 3ZnTPP* and methyl viologen in lecithin vesicles studied by global analysis. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> . 1994. 82:149-160
Francesca Cutruzzol. Bacterial nitric oxide synthesis. <i>Biochimica et Biophysica Acta (BBA) - Bioenergetics</i> . 1999. 1411:231-249
Guozheng Deblois Dai. Differential sensitivity of five cyanobacterial strains to ammonium toxicity and its inhibitory mechanism on the photosynthesis of rice-field cyanobacterium <i>Ge. Aquatic Toxicology</i> . 2008. 89:113-121
Antonio L. Santamaria De Lacey. Kinetic characterization of <i>Desulfovibrio gigas</i> hydrogenase upon selective chemical modification of amino acid groups as a tool for structure. <i>Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology</i> . 2000. 1481:371-380
Deepika Miller de Silva. In vitro loading of apoferritin. <i>Archives of Biochemistry and Biophysics</i> . 1992. 293:409-415
Chantal Roullier Degrand. Quinonoid modified electrodes amplify the current from certain soluble one-electron redox reagents. <i>Journal of Electroanalytical Chemistry and Interfacial Electrochemistry</i> . 1984. 178:101-111
Janice A. Rao DeGray. Reduction of paraquat and related bipyridylum compounds to free radical metabolites by rat hepatocytes. <i>Archives of Biochemistry and Biophysics</i> . 1991. 289:145-152
Anissa Adhoum Dhaouadi. Degradation of paraquat herbicide by electrochemical advanced oxidation methods. <i>Journal of Electroanalytical Chemistry</i> . 2009. 637:33-42
Anissa Adhoum Dhaouadi. Heterogeneous catalytic wet peroxide oxidation of paraquat in the presence of modified activated carbon. <i>Applied Catalysis B: Environmental</i> . 2010. 97:227-235
Richard A. Dilley. The effect of various energy-conversion states of chloroplasts on proton and electron transport. <i>Archives of Biochemistry and Biophysics</i> . 1970. 137:270-283
Ricardo Jorge de Pinho Dinis-Oliveira. Reactivity of paraquat with sodium salicylate: Formation of stable complexes. <i>Toxicology</i> . 2008. 249:130-139
J. Schaap Klimek. Effect of paraquat on cytochrome P-450-dependent lipid peroxidation in bovine adrenal cortex mitochondria. <i>Biochim Biophys Acta</i> . 1983. 752:127-36
Matthias Klemme Drutschmann. Sulfide-repressed, membrane-bound hydrogenase in the thermophilic facultative phototroph, <i>Chloroflexus aurantiacus</i> . <i>FEMS Microbiology Letters</i> . 1985. 28:231-235
Euan Ip Duncanson. Synthesis of nitrite reductase is regulated differently in leaf and root of barley (<i>Hordeum vulgare</i> L.). <i>Plant Science</i> . 1992. 87:151-160
Tumpa Das Dutta. Purification and characterization of [Fe]-hydrogenase from high yielding hydrogen-producing strain, <i>Enterobacter cloacae</i> IIT-BT08 (MTCC 5373). <i>International Journal of Hydrogen Energy</i> . 2009. 34:7530-7537
Takeshi Ageishi Endo. HYdrogen Evolution by use of 1-alkyl-1,4-dihydronicotinamide derivatives. <i>Journal of Molecular Catalysis</i> . 1983. 18:351-358

Chlorophyll fluorescence measurements were performed on osmotically lysed potato chloroplasts in order to characterize
The immobilization of nitrate reductase (NR) was performed by entrapment in a laponite clay gel and cross-linking by glutaraldehyde
The effects were studied of the plastoquinone analogs 2,5-dibromo-3-methyl-6-isopropyl-p-benzoquinone (DBMIB) and tetrachloro-2,5-dimethyl-p-benzoquinone (TCMQ)
Suspensions of lecithin vesicles incorporating zinc tetraphenylporphyrin in high and low local concentrations (lipid-to-porphyrin ratio)
The structure-function relationships in nitrite reductases, key enzymes in the dissimilatory denitrification pathway which
Effects of two fertilizers, NH ₄ Cl and KCl, on the growth of the edible cyanobacterium <i>Geotrichum</i>
The effect of amino acid residues modification of <i>Desulfovibrio gigas</i> hydrogenase on different activity assays is reported
This study compared the effect of loading apoferritin either with ferrous ammonium sulfate in various buffers or with cerium(IV) ammonium nitrate
Chemically modified carbon electrodes are prepared which hold polymeric layers of anthraquinone or dopamine units on their surface
The toxicity of paraquat is due to the oxygen-derived radicals formed by the reaction of oxygen with bipyridylum radical
Various electrochemical advanced oxidation processes (EAOPs) including anodic oxidation (AO), electro-Fenton (EF) and electro- <i>peroxone</i> (EP)
The catalytic wet peroxide oxidation (CWPO) method was applied to the degradation of paraquat, a widely used and highly toxic herbicide
The stoichiometry of H ⁺ and electron transport in spinach chloroplasts was very sensitive to the presence or absence of thylakoid membranes
Sodium salicylate (NaSAL) has been shown to be a promising antidote for the treatment of paraquat (PQ) poisonings. The
We have investigated the effect of paraquat (methyl viologen) on lipid peroxidation in bovine adrenal cortex mitochondria
The thermophilic facultatively phototrophic green bacterium <i>Chloroflexus aurantiacus</i> strain Ok-70-fl was shown to possess
Activity staining after non-denaturing polyacrylamide gel electrophoresis (PAGE) of extracts from nitrate-treated plants indicated
Fe-hydrogenase from <i>Enterobacter cloacae</i> IIT-BT08 was purified 1284 fold with specific activity of 335 μ mol H ₂ /min/mg
Summary It was found that hydrogen was produced from 1,4-dihydronicotinamide derivatives such as 1-benzyl-1,4-dihydronicotinamide

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant